

AMENDMENT TO THE SPECIFICATION

Please amend the specification as follows, without prejudice or disclaimer.

Please replace paragraph [0018] with the following amended paragraph:

[0018] TAs are typically classified into five categories according to their expression pattern, function, or genetic origin: cancer-testis (CT) antigens (i.e., MAGE, NY-ESO-1); melanocyte differentiation antigens (i.e., Melan A/MART-1, tyrosinase, gp100); mutational antigens (i.e., MUM-1, p53, CDK-4); overexpressed 'self' antigens (i.e., HER-2/neu, p53); and, viral antigens (i.e., HPV, EBV). For the purposes of practicing the present invention, a suitable TA is any TA that induces or enhances an anti-tumor immune response in a host to whom the TA has been administered. Suitable TAs include, for example, gp100 (Cox et al., *Science*, 264:716-719 (1994)), MART-1/Melan. A (Kawakami et al., *J. Exp. Med.*, 180:347-352 (1994)), gp75 (TRP-1) (Wang et al., *J. Exp. Med.*, 186:1131-1140 (1996)), tyrosinase (Wolfel et al., *Eur. J Immunol.*, 24:759-764 (1994); WO 200175117; WO 200175016; WO 200175007), NY-ESO-1 (WO 98/14464 (U.S. Pat. No. 6,274,145B1); WO 99/18206), melanoma proteoglycan (Hellstrom et al., *J. Immunol.*, 130:1467-1472 (1983)), MAGE family antigens (i.e., MAGE-1, 2,3,4,6,12, 51; Van der Bruggen et al., *Science*, 254:1643-1647 (1991); U.S. Pat. No. 6,235,525; CN 1319611), BAGE family antigens (Boel et al., *Immunity*, 2:167-175 (1995), U.S. Pat. No. 5,571,711), GAGE family antigens (i.e., GAGE-1,2; Van den Eynde et al., *J. Exp. Med.*, 182:689-698 (1995), U.S. Pat. No. 5,610,013); DAGE family antigens (U.S. Pat. No., 6,013,765); RAGE family antigens (i.e., RAGE-1; Gaugler et al., *Immunogenetics*, 44:323-330 (1996); U.S. Pat. No. 5,939,526), N-acetylglucosaminyltransferase-V (Guilloux et al., *J. Exp. Med.*, 183:1173-1183 (1996)), p15 (Robbins et al., *J. Immunol.* 154:5944-5950 (1995)), β -catenin (Robbins et al., *J. Exp. Med.*, 183:1185-1192 (1996)), MUM-1 (Coulie et al., *Proc. Natl. Acad. Sci. USA*, 92:7976-7980 (1995)), cyclin dependent kinase-4 (CDK4) (Wolfel et al., *Science*, 269:1281-1284 (1995)), p21-ras (Fossum et al., *Int. J. Cancer*, 56:40-45 (1994)), BCR-abl (Bocchia et al., *Blood*, 85:2680-2684 (1995)), p53 (Theobald et al., *Proc. Natl. Acad. Sci. USA*, 92:11993-11997 (1995)), p185 HER2/neu (erb-B1; Fisk et al., *J. Exp. Med.*, 181:2109-2117 (1995)), epidermal growth factor receptor (EGFR) (Harris et al., *Breast Cancer Res. Treat*, 29:1-2

(1994)), carcinoembryonic antigens (CEA) (Kwong et al., *J. Natl. Cancer Inst.*, 85:982-990 (1995) U.S. Pat. Nos. 5,756,103; 5,274,087; 5,571,710; 6,071,716; 5,698,530; 6,045,802; EP 263933; EP 346710; and, EP 784483); carcinoma-associated mutated mucins (i.e., MUC-1 gene products; Jerome et al., *J. Immunol.*, 151:1654-1662 (1993)); EBNA gene products of EBV (i.e., EBNA-1; Rickinson et al., *Cancer Surveys*, 13:53-80 (1992)); E7, E6 proteins of human papillomavirus (Ressing et al., *J. Immunol.*, 154:5934-5943 (1995)); prostate specific antigen (PSA; Xue et al., *The Prostate*, 30:73-78 (1997)); prostate specific membrane antigen (PSMA; Israeli, et al., *Cancer Res.*, 54:1807-1811 (1994)); idiotypic epitopes or antigens, for example, immunoglobulin idiotypes or T cell receptor idiotypes (Chen et al., *J. Immunol.*, 153:4775-4787 (1994)); KSA (U.S. Pat. No. 5,348,887), kinesin 2 (Dietz, et al. *Biochem Biophys Res Commun* 2000 Sep 7;275(3):731-8), HIP-55, TGF β -1 anti-apoptotic factor (Toomey, et al. *Br J Biomed Sci* 2001;58(3):177-83), tumor protein D52 (Bryne J. A., et al., *Genomics*, 35:523-532 (1996)), H1FT, NY-BR-1 (WO 01/47959), NY-BR-62, NY-BR-75, NY-BR-85, NY-BR-87, NY-BR-96 (Scanlan, M. *Serologic and Bioinformatic Approaches to the Identification of Human Tumor Antigens*, in *Cancer Vaccines 2000*, Cancer Research Institute, New York, N.Y.), including "wild-type" (i.e., normally encoded by the genome, naturally-occurring), modified, and mutated versions as well as other fragments and derivatives thereof. Any of these TAs may be utilized alone or in combination with one another in a co-immunization protocol.